


CS-GY 6763: Lecture 12

Stochastic Block Model, Randomized numerical linear algebra, ϵ -net arguments.

NYU Tandon School of Engineering, Prof. Christopher Musco

LAST CLASS

Represent undirected graph as symmetric matrix: $n \times n$
adjacency matrix A and graph Laplacian $L = D - A$ where D is
the diagonal degree matrix.



$$\begin{matrix} & \mathbf{D} & & \mathbf{A} & & \mathbf{L} \end{matrix}$$
$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 3 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 2 \end{bmatrix} - \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & -1 & 0 & 0 \\ -1 & 3 & -1 & -1 \\ 0 & -1 & 2 & -1 \\ 0 & -1 & -1 & 2 \end{bmatrix}$$

$L = B^T B$ where B is the “edge-vertex incidence” matrix.

$$B = \begin{bmatrix} 1 & -1 & 0 & 0 \\ 0 & 1 & -1 & 0 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & -1 \end{bmatrix}$$

Balanced Cut: Partition nodes along a cut that:

- Has few crossing edges: $|\{(u, v) \in E : u \in S, v \in T\}|$ is small.
- Separates large partitions: $|S|, |T|$ are not too small.



(a) Zachary Karate Club Graph

We observed that $\mathbf{x}^T L \mathbf{x} = \sum_{(i,j) \in E} (\mathbf{x}(i) - \mathbf{x}(j))^2$. If \mathbf{c} is a “cut indicator vector” for a cut between node set S and T – i.e. $\mathbf{c}[i] = 1$ for all $i \in S$ and -1 for $i \in T$, then it followed that:

$$\mathbf{c}^T L \mathbf{c} = 4 \cdot \text{cut}(S, T).$$

Note: \mathbf{c} often denote by $\chi_{S,T}$ to remind us what the cut is. And recall that we always have $S = V \setminus T$.

“Relax and round” algorithm:

- Relax problem $\min \mathbf{c}^T \mathbf{L} \mathbf{c}$ by not requiring \mathbf{c} to be a binary cut-indicator vector.
- Showed that second smallest eigenvector \mathbf{v}_{n-1} of \mathbf{L} solved the relaxed “perfectly balanced” cut problem.
- Round this vector to be a cut indicator vector: all negative entries rounded to -1 , all positive entries rounded to 1 .

Main theoretical result: This approach is hard to analyze in general, but can be proven to work well on random graphs drawn from a natural generative model!.

GENERATIVE MODELS

So far: Showed that spectral clustering partitions a graph along a small cut between large pieces.

- No formal guarantee on the ‘quality’ of the partitioning.
- Difficult to analyze for general input graphs.

Common approach: Design a natural **generative model** that produces random but realistic inputs and analyze how the algorithm performs on inputs drawn from this model.

- Very common in algorithm design and analysis. Great way to start approaching a problem.
- This is also the whole idea behind Bayesian Machine Learning (can be used to justify ℓ_2 linear regression, k -means clustering, PCA, etc.)


Ideas for a generative model for **social network graphs** that would allow us to understand partitioning?

STOCHASTIC BLOCK MODEL

Stochastic Block Model (Planted Partition Model):

Let $G_n(p, q)$ be a distribution over graphs on n nodes, split equally into two groups B and C , each with $n/2$ nodes.


- Any two nodes in the **same group** are connected with probability p (including self-loops).
- Any two nodes in **different groups** are connected with prob. $q < p$.



LINEAR ALGEBRAIC VIEW

Let G be a stochastic block model graph drawn from $G_n(p, q)$.


- Let $A \in \mathbb{R}^{n \times n}$ be the adjacency matrix of G . What is $\mathbb{E}[A]$?



Note that we are arbitrarily ordering the nodes in A by group. In reality A would look “scrambled” as on the right.

STOCHASTIC BLOCK MODEL

Goal is to find the “ground truth” balanced partition B, C using our standard spectral method.




To do so, we need to understand the second smallest eigenvalue of $\mathbf{L} = \mathbf{D} - \mathbf{A}$. We will start by considering the expected value of these matrices:

$$\mathbb{E}[\mathbf{L}] = \mathbb{E}[\mathbf{D}] - \mathbb{E}[\mathbf{A}].$$

EXPECTED ADJACENCY SPECTRUM

Letting G be a stochastic block model graph drawn from $G_n(p, q)$ and $\mathbf{A} \in \mathbb{R}^{n \times n}$ be its adjacency matrix. $(\mathbb{E}[\mathbf{A}])_{i,j} = p$ for i, j in same group, $(\mathbb{E}[\mathbf{A}])_{i,j} = q$ otherwise.




We are going to determine the eigenvectors and eigenvalues of $\mathbb{E}[\mathbf{A}]$.

What is the expected Laplacian of $G_n(p, q)$?

$\mathbb{E}[\mathbf{A}]$ and $\mathbb{E}[\mathbf{L}]$ have the same eigenvectors and eigenvalues are equal up to a shift/inversion. So second largest eigenvector of $\mathbb{E}[\mathbf{A}]$ is the same as the second smallest of $\mathbb{E}[\mathbf{L}]$

EXPECTED ADJACENCY SPECTRUM

Letting G be a stochastic block model graph drawn from $G_n(p, q)$ and $\mathbf{A} \in \mathbb{R}^{n \times n}$ be its adjacency matrix, what are the eigenvectors and eigenvalues of $\mathbb{E}[\mathbf{A}]$?



EXPECTED ADJACENCY SPECTRUM

Diagram illustrating the Expected Adjacency Spectrum. The matrix $E[A]$ is a 2×2 block matrix with blocks p and q , representing communities B and C (each with $n/2$ nodes). The matrix is decomposed into $V \Lambda V^T$.

The matrix $E[A]$ is shown as a 2×2 block matrix with blocks p and q , representing communities B and C (each with $n/2$ nodes). The matrix is decomposed into $V \Lambda V^T$.

The matrix V is a 2×8 matrix with columns \mathbf{v}_1 and \mathbf{v}_2 .

The matrix Λ is a diagonal matrix with eigenvalues $\frac{p+q}{2}$ and $\frac{p-q}{2}$.

The matrix V^T is the transpose of V .

- $\mathbf{v}_1 \sim \mathbf{1}$ with eigenvalue $\lambda_1 = \frac{(p+q)n}{2}$.
- $\mathbf{v}_2 \sim \chi_{B,C}$ with eigenvalue $\lambda_2 = \frac{(p-q)n}{2}$.

If we compute \mathbf{v}_2 then we recover the communities B and C !

Upshot: The second smallest eigenvector of $\mathbb{E}[\mathbf{L}]$, equivalently the second largest of $\mathbb{E}[\mathbf{A}]$, is exactly $\chi_{B,C}$ – the indicator vector for the cut between the communities.

- If the random graph G (equivalently \mathbf{A} and \mathbf{L}) were exactly equal to its expectation, partitioning using this eigenvector would exactly recover communities B and C .

How do we show that a matrix (e.g., \mathbf{A}) is close to its expectation? **Matrix concentration inequalities.**

- Analogous to scalar concentration inequalities like Markovs, Chebyshevs, Bernsteins.

MATRIX CONCENTRATION

Alon, Krivelevich, Vu, 2002:

Matrix Concentration Inequality: If $p \geq O\left(\frac{\log^4 n}{n}\right)$, then with high probability

$$\|\mathbf{A} - \mathbb{E}[\mathbf{A}]\|_2 \leq O(\sqrt{pn}).$$

where $\|\cdot\|_2$ is the matrix **spectral** norm (operator norm).

Recall that $\|\mathbf{X}\|_2 = \max_{\mathbf{z} \in \mathbb{R}^d: \|\mathbf{z}\|_2=1} \|\mathbf{X}\mathbf{z}\|_2 = \sigma_1(\mathbf{X})$.

$\|\mathbf{A}\|_2$ is on the order of $O(p\sqrt{n})$ so another way of thinking about the right hand side is $\frac{\|\mathbf{A}\|_2}{\sqrt{p}}$. I.e. get's better with p .

EIGENVECTOR PERTURBATION

For the stochastic block model application, we want to show that the second eigenvectors of \mathbf{A} and $\mathbb{E}[\mathbf{A}]$ are close. How does this relate to their difference in spectral norm?

Davis-Kahan Eigenvector Perturbation Theorem: Suppose $\mathbf{A}, \bar{\mathbf{A}} \in \mathbb{R}^{d \times d}$ are symmetric with $\|\mathbf{A} - \bar{\mathbf{A}}\|_2 \leq \epsilon$ and eigenvectors $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ and $\bar{\mathbf{v}}_1, \bar{\mathbf{v}}_2, \dots, \bar{\mathbf{v}}_n$. Letting $\theta(\mathbf{v}_i, \bar{\mathbf{v}}_i)$ denote the angle between \mathbf{v}_i and $\bar{\mathbf{v}}_i$, for all i :

$$\sin[\theta(\mathbf{v}_i, \bar{\mathbf{v}}_i)] \leq \frac{\epsilon}{\min_{j \neq i} |\lambda_i - \lambda_j|}$$

where $\lambda_1, \dots, \lambda_n$ are the eigenvalues of $\bar{\mathbf{A}}$.

APPLICATION TO STOCHASTIC BLOCK MODEL

Claim 1 (Matrix Concentration): For $p \geq O\left(\frac{\log^4 n}{n}\right)$,

$$\|A - \mathbb{E}[A]\|_2 \leq O(\sqrt{pn}).$$

Claim 2 (Davis-Kahan): For $p \geq O\left(\frac{\log^4 n}{n}\right)$,

$$\sin \theta(\mathbf{v}_2, \bar{\mathbf{v}}_2) \leq \frac{O(\sqrt{pn})}{\min_{j \neq i} |\lambda_i - \lambda_j|} \leq \frac{O(\sqrt{pn})}{(p-q)n/2} = O\left(\frac{\sqrt{p}}{(p-q)\sqrt{n}}\right)$$

Recall: $\mathbb{E}[A]$, has eigenvalues $\lambda_1 = \frac{(p+q)n}{2}$, $\lambda_2 = \frac{(p-q)n}{2}$, $\lambda_i = 0$ for $i \geq 3$.

$$\min_{j \neq i} |\lambda_i - \lambda_j| = \min \left(qn, \frac{(p-q)n}{2} \right).$$


Assume $\frac{(p-q)n}{2}$ will be the minimum of these two gaps.

(A slightly trickier analysis can remove the qn term entirely.)


APPLICATION TO STOCHASTIC BLOCK MODEL

So far: $\sin \theta(\mathbf{v}_2, \bar{\mathbf{v}}_2) \leq O\left(\frac{\sqrt{p}}{(p-q)\sqrt{n}}\right)$. What does this give us?

- Can show that this implies $\|\mathbf{v}_2 - \bar{\mathbf{v}}_2\|_2^2 \leq O\left(\frac{p}{(p-q)^2 n}\right)$ (exercise).
- $\bar{\mathbf{v}}_2$ is $\frac{1}{\sqrt{n}}\chi_{B,C}$: the community indicator vector.



- To understand how well rounding recovers $\bar{\mathbf{v}}_2$, need to understand how many locations \mathbf{v}_2 and $\bar{\mathbf{v}}_2$ can differ in sign.



Main argument:

- Every i where $v_2(i), \bar{v}_2(i)$ differ in sign contributes $\geq \frac{1}{n}$ to $\|\mathbf{v}_2 - \bar{\mathbf{v}}_2\|_2^2$.
- We know that $\|\mathbf{v}_2 - \bar{\mathbf{v}}_2\|_2^2 \leq O\left(\frac{p}{(p-q)^2 n}\right)$.
- So \mathbf{v}_2 and $\bar{\mathbf{v}}_2$ differ in sign in at most $O\left(\frac{p}{(p-q)^2}\right)$ positions.

Upshot: If G is a stochastic block model graph with adjacency matrix \mathbf{A} , if we compute its second large eigenvector \mathbf{v}_2 and assign nodes to communities according to the sign pattern of this vector, we will correctly assign all but $O\left(\frac{p}{(p-q)^2}\right)$ nodes.

- **Hard case:** $p = c/n$ for some factor c . Even when $p - q = O(1/n)$, assign all but an $O(n)$ fraction of nodes correctly. E.g., assign 99% of nodes to the right cluster.

Forget about the previous problem, but still consider the matrix $\mathbf{M} = \mathbb{E}[\mathbf{A}]$.


- Dense $n \times n$ matrix.
- Computing top eigenvectors takes $\approx O(n^2/\sqrt{\epsilon})$ time.

If someone asked you to speed this up and return approximate top eigenvectors, what could you do?


RANDOMIZED NUMERICAL LINEAR ALGEBRA

Main idea: If you want to compute singular vectors, multiply two matrices, solve a regression problem, etc.:


1. Compress your matrices using a randomized method (e.g. subsampling).
2. Solve the problem on the smaller or sparser matrix.
 - \tilde{A} called a “sketch” or “coreset” for A .



Approximate matrix multiplication:




Approximate regression:



SKETCHED REGRESSION

Today's example: Randomized approximate regression using a Johnson-Lindenstrauss Matrix.



Input: $A \in \mathbb{R}^{n \times d}$, $b \in \mathbb{R}^n$.

Goal: Let $x^* = \arg \min_x \|Ax - b\|_2^2$. Let $\tilde{x} = \arg \min_x \|\Pi Ax - \Pi b\|_2^2$

Want: $\|A\tilde{x} - b\|_2^2 \leq (1 + \epsilon) \|Ax^* - b\|_2^2$

Theorem (Randomized Linear Regression)

Let Π be a properly scaled JL matrix (random Gaussian, sign, sparse random, etc.) with $m = O\left(\frac{d}{\epsilon^2}\right)$ rows¹. Then with probability 9/10, for any $\mathbf{A} \in \mathbb{R}^{n \times d}$ and $\mathbf{b} \in \mathbb{R}^n$,

$$\|\mathbf{A}\tilde{\mathbf{x}} - \mathbf{b}\|_2^2 \leq (1 + \epsilon)\|\mathbf{A}\mathbf{x}^* - \mathbf{b}\|_2^2$$

where $\tilde{\mathbf{x}} = \arg \min_{\mathbf{x}} \|\Pi \mathbf{A} \mathbf{x} - \Pi \mathbf{b}\|_2^2$.

¹This can be improved to $O(d/\epsilon)$ with a tighter analysis

- Prove this theorem using an ϵ -net argument, which is a popular technique for applying our standard concentration inequality + union bound argument to an infinite number of events.
- These sort of arguments appear all the time in theoretical algorithms and ML research, so this part of lecture is as much about the technique as the final result.
- You will use an ϵ -net argument to prove a matrix concentration inequality on your last problem set.

Claim: Suffices to prove that for all $\mathbf{x} \in \mathbb{R}^d$,

$$(1 - \epsilon)\|\mathbf{Ax} - \mathbf{b}\|_2^2 \leq \|\mathbf{\Pi Ax} - \mathbf{\Pi b}\|_2^2 \leq (1 + \epsilon)\|\mathbf{Ax} - \mathbf{b}\|_2^2$$

Lemma (Distributional JL)

If $\mathbf{\Pi}$ is chosen to a properly scaled random Gaussian matrix, sign matrix, sparse random matrix, etc., with $O\left(\frac{\log(1/\delta)}{\epsilon^2}\right)$ rows then for any fixed \mathbf{y} ,

$$(1 - \epsilon)\|\mathbf{y}\|_2^2 \leq \|\mathbf{\Pi y}\|_2^2 \leq (1 + \epsilon)\|\mathbf{y}\|_2^2$$

with probability $(1 - \delta)$.

Corollary: For any fixed \mathbf{x} , with probability $(1 - \delta)$,

$$(1 - \epsilon)\|\mathbf{Ax} - \mathbf{b}\|_2^2 \leq \|\mathbf{\Pi Ax} - \mathbf{\Pi b}\|_2^2 \leq (1 + \epsilon)\|\mathbf{Ax} - \mathbf{b}\|_2^2.$$

How do we go from “for any fixed \mathbf{x} ” to “for all $\mathbf{x} \in \mathbb{R}^d$ ”.

This statement requires establishing a Johnson-Lindenstrauss type bound for an infinity of possible vectors $(\mathbf{Ax} - \mathbf{b})$, which can't be tackled directly with a union bound argument.


Note that all vectors of the form $(\mathbf{Ax} - \mathbf{b})$ lie in a low dimensional subspace: spanned by $d + 1$ vectors, where d is the width of \mathbf{A} . **So even though the set is infinite, it is “simple” in some way. Parameterized by just $d + 1$ numbers.**

Theorem (Subspace Embedding from JL)

Let $\mathcal{U} \subset \mathbb{R}^n$ be a d -dimensional linear subspace in \mathbb{R}^n . If $\mathbf{\Pi} \in \mathbb{R}^{m \times d}$ is chosen from any distribution \mathcal{D} satisfying the Distributional JL Lemma, then with probability $1 - \delta$,

$$(1 - \epsilon)\|\mathbf{v}\|_2^2 \leq \|\mathbf{\Pi v}\|_2^2 \leq (1 + \epsilon)\|\mathbf{v}\|_2^2$$

for all $\mathbf{v} \in \mathcal{U}$, as long as $m = O\left(\frac{d \log(1/\epsilon) + \log(1/\delta)}{\epsilon^2}\right)$.²



²It's possible to obtain a slightly tighter bound of $O\left(\frac{d + \log(1/\delta)}{\epsilon^2}\right)$. It's a nice challenge to try proving this.

Corollary: If we choose Π and properly scale, then with $O(d/\epsilon^2)$ rows,

$$(1 - \epsilon) \| \mathbf{Ax} - \mathbf{b} \|_2^2 \leq \| \Pi \mathbf{Ax} - \Pi \mathbf{b} \|_2^2 \leq (1 + \epsilon) \| \mathbf{Ax} - \mathbf{b} \|_2^2$$

for all \mathbf{x} and thus

$$\| \mathbf{A}\tilde{\mathbf{x}} - \mathbf{b} \|_2^2 \leq (1 + O(\epsilon)) \min_{\mathbf{x}} \| \mathbf{Ax} - \mathbf{b} \|_2^2.$$

I.e., our main theorem is proven.


Proof: Apply Subspace Embedding Thm. to the $(d + 1)$ dimensional subspace spanned by \mathbf{A} 's d columns and \mathbf{b} . Every vector $\mathbf{Ax} - \mathbf{b}$ lies in this subspace.

Theorem (Subspace Embedding from JL)

Let $\mathcal{U} \subset \mathbb{R}^n$ be a d -dimensional linear subspace in \mathbb{R}^n . If $\mathbf{\Pi} \in \mathbb{R}^{m \times d}$ is chosen from any distribution \mathcal{D} satisfying the Distributional JL Lemma, then with probability $1 - \delta$,

$$(1 - \epsilon)\|\mathbf{v}\|_2^2 \leq \|\mathbf{\Pi v}\|_2^2 \leq (1 + \epsilon)\|\mathbf{v}\|_2^2 \quad (1)$$

for all $\mathbf{v} \in \mathcal{U}$, as long as $m = O\left(\frac{d \log(1/\epsilon) + \log(1/\delta)}{\epsilon^2}\right)$



Subspace embeddings have tons of other applications!

$$(1 - \epsilon)\|\mathbf{v}\|_2^2 \leq \|\Pi\mathbf{v}\|_2^2 \leq (1 + \epsilon)\|\mathbf{v}\|_2^2 \quad (2)$$

First Observation: The theorem holds as long as (2) holds for all \mathbf{w} on the unit sphere in \mathcal{U} . Denote the sphere $S_{\mathcal{U}}$:


$$S_{\mathcal{U}} = \{\mathbf{w} \mid \mathbf{w} \in \mathcal{U} \text{ and } \|\mathbf{w}\|_2 = 1\}.$$

Follows from linearity: Any point $\mathbf{v} \in \mathcal{U}$ can be written as $c\mathbf{w}$ for some scalar c and some point $\mathbf{w} \in S_{\mathcal{U}}$.

- If $(1 - \epsilon)\|\mathbf{w}\|_2 \leq \|\Pi\mathbf{w}\|_2 \leq (1 + \epsilon)\|\mathbf{w}\|_2$.
- then $c(1 - \epsilon)\|\mathbf{w}\|_2 \leq c\|\Pi\mathbf{w}\|_2 \leq c(1 + \epsilon)\|\mathbf{w}\|_2$,
- and thus $(1 - \epsilon)\|c\mathbf{w}\|_2 \leq \|\Pi c\mathbf{w}\|_2 \leq (1 + \epsilon)\|c\mathbf{w}\|_2$.

SUBSPACE EMBEDDING PROOF

Intuition: There are not too many “different” points on a d -dimensional sphere:



N_ϵ is called an “ ϵ ”-net.

If we can prove

$$(1 - \epsilon)\|\mathbf{w}\|_2 \leq \|\Pi\mathbf{w}\|_2 \leq (1 + \epsilon)\|\mathbf{w}\|_2$$

for all points $\mathbf{w} \in N_\epsilon$, we can hopefully extend to all of S_U .

Lemma (ϵ -net for the sphere)

For any $\epsilon \leq 1$, there exists a set $N_\epsilon \subset S_{\mathcal{U}}$ with $|N_\epsilon| = \left(\frac{4}{\epsilon}\right)^d$ such that $\forall \mathbf{v} \in S_{\mathcal{U}}$,

$$\min_{\mathbf{w} \in N_\epsilon} \|\mathbf{v} - \mathbf{w}\|_2 \leq \epsilon.$$

Take this claim to be true for now: we will prove later.

1. Preserving norms of all points in net N_ϵ .

Set $\delta' = \frac{1}{|N_\epsilon|} \cdot \delta = \left(\frac{\epsilon}{4}\right)^d \cdot \delta$. As long as Π has $O\left(\frac{\log(1/\delta')}{\epsilon^2}\right) = O\left(\frac{d \log(1/\epsilon) + \log(1/\delta)}{\epsilon^2}\right)$ rows, then by a union bound,

$$(1 - \epsilon)\|\mathbf{w}\|_2 \leq \|\Pi\mathbf{w}\|_2 \leq (1 + \epsilon)\|\mathbf{w}\|_2.$$


for all $\mathbf{w} \in N_\epsilon$, with probability $1 - \delta$.

2. Writing any point in sphere as linear comb. of points in N_ϵ .

For some $w_0, w_1, w_2 \dots \in N_\epsilon$, any $v \in S_{\mathcal{U}}$ can be written:

$$v = w_0 + c_1 w_1 + c_2 w_2 + \dots$$

for constants c_1, c_2, \dots where $|c_i| \leq \epsilon^i$.



3. Preserving norm of v .

Applying triangle inequality, we have that:

$$\begin{aligned}
 \|\Pi v\|_2 &= \|\Pi w_0 + c_1 \Pi w_1 + c_2 \Pi w_2 + \dots\| \\
 &\leq \|\Pi w_0\| + c_1 \|\Pi w_1\| + c_2 \|\Pi w_2\| + \dots \\
 &\leq \|\Pi w_0\| + \epsilon \|\Pi w_1\| + \epsilon^2 \|\Pi w_2\| + \dots \\
 &\leq (1 + \epsilon) + \epsilon(1 + \epsilon) + \epsilon^2(1 + \epsilon) + \dots \\
 &\leq 1 + 2\epsilon.
 \end{aligned}$$

3. Preserving norm of v .

Similarly,

$$\begin{aligned}
 \|\Pi v\|_2 &= \|\Pi w_0 + c_1 \Pi w_1 + c_2 \Pi w_2 + \dots\| \\
 &\geq \|\Pi w_0\| - \epsilon \|\Pi w_1\| - \epsilon^2 \|\Pi w_2\| - \dots \\
 &\geq (1 - \epsilon) - \epsilon(1 + \epsilon) - \epsilon^2(1 + \epsilon) - \dots \\
 &\geq 1 - 5\epsilon.
 \end{aligned}$$

So we have proven

$$(1 - O(\epsilon)) \|\mathbf{v}\|_2 \leq \|\mathbf{\Pi v}\|_2 \leq (1 + O(\epsilon)) \|\mathbf{v}\|_2$$

for all $\mathbf{v} \in S_{\mathcal{U}}$, which in turn implies,

$$(1 - O(\epsilon)) \|\mathbf{v}\|_2^2 \leq \|\mathbf{\Pi v}\|_2^2 \leq (1 + O(\epsilon)) \|\mathbf{v}\|_2^2$$

Adjusting ϵ proves the Subspace Embedding theorem.

Theorem (Subspace Embedding from JL)

Let $\mathcal{U} \subset \mathbb{R}^n$ be a d -dimensional linear subspace in \mathbb{R}^n . If $\mathbf{\Pi} \in \mathbb{R}^{m \times d}$ is chosen from any distribution \mathcal{D} satisfying the Distributional JL Lemma, then with probability $1 - \delta$,

$$(1 - \epsilon)\|\mathbf{v}\|_2^2 \leq \|\mathbf{\Pi v}\|_2^2 \leq (1 + \epsilon)\|\mathbf{v}\|_2^2 \quad (3)$$

for all $\mathbf{v} \in \mathcal{U}$, as long as $m = O\left(\frac{d \log(1/\epsilon) + \log(1/\delta)}{\epsilon^2}\right)$

Subspace embeddings have many other applications!

For example, if $m = O(k/\epsilon)$, $\mathbf{\Pi A}$ can be used to compute an approximate partial SVD, which leads to a $(1 + \epsilon)$ approximate low-rank approximation for \mathbf{A} .

Lemma (ϵ -net for the sphere)

For any $\epsilon \leq 1$, there exists a set $N_\epsilon \subset S_{\mathcal{U}}$ with $|N_\epsilon| = \left(\frac{3}{\epsilon}\right)^d$ such that $\forall \mathbf{v} \in S_{\mathcal{U}}$,


$$\min_{\mathbf{w} \in N_\epsilon} \|\mathbf{v} - \mathbf{w}\| \leq \epsilon.$$

Imaginary algorithm for constructing N_ϵ :

- Set $N_\epsilon = \{\}$
- While such a point exists, choose an arbitrary point $\mathbf{v} \in S_{\mathcal{U}}$ where $\nexists \mathbf{w} \in N_\epsilon$ with $\|\mathbf{v} - \mathbf{w}\| \leq \epsilon$. Set $N_\epsilon = N_\epsilon \cup \{\mathbf{w}\}$.

After running this procedure, we have $N_\epsilon = \{\mathbf{w}_1, \dots, \mathbf{w}_{|N_\epsilon|}\}$ and $\min_{\mathbf{w} \in N_\epsilon} \|\mathbf{v} - \mathbf{w}\| \leq \epsilon$ for all $\mathbf{v} \in S_{\mathcal{U}}$ as desired.

How many steps does this procedure take?



Can place a ball of radius $\epsilon/2$ around each w_i without intersecting any other balls. All of these balls live in a ball of radius $1 + \epsilon/2$.

Volume of d dimensional ball of radius r is

$$\text{vol}(d, r) = c \cdot r^d,$$

where c is a constant that depends on d , but not r . From previous slide we have:

$$\begin{aligned}\text{vol}(d, \epsilon/2) \cdot |N_\epsilon| &\leq \text{vol}(d, 1 + \epsilon/2) \\ |N_\epsilon| &\leq \frac{\text{vol}(d, 1 + \epsilon/2)}{\text{vol}(d, \epsilon/2)} \\ &\leq \left(\frac{1 + \epsilon/2}{\epsilon/2} \right)^d \leq \left(\frac{3}{\epsilon} \right)^d\end{aligned}$$

You can actually show that $m = O\left(\frac{d + \log(1/\delta)}{\epsilon}\right)$ suffices to be a d dimensional subspace embedding, instead of the bound we proved of $m = O\left(\frac{d \log(1/\epsilon) + \log(1/\delta)}{\epsilon}\right)$.

The trick is to show that a constant factor net is actually all that you need instead of an ϵ factor.

RUNTIME CONSIDERATION


For $\epsilon, \delta = O(1)$, we need $\mathbf{\Pi}$ to have $m = O(d)$ rows.

- Cost to solve $\|\mathbf{Ax} - \mathbf{b}\|_2^2$:
 - $O(nd^2)$ time for direct method. Need to compute $(\mathbf{A}^T \mathbf{A})^{-1} \mathbf{A}^T \mathbf{b}$.
 - $O(nd) \cdot (\text{\# of iterations})$ time for iterative method (GD, AGD, conjugate gradient method).
- Cost to solve $\|\mathbf{\Pi Ax} - \mathbf{\Pi b}\|_2^2$:
 - $O(d^3)$ time for direct method.
 - $O(d^2) \cdot (\text{\# of iterations})$ time for iterative method.

RUNTIME CONSIDERATION

But time to compute ΠA is an $(m \times n) \times (n \times d)$ matrix multiply: $O(mnd) = O(nd^2)$ time!

Goal: Develop faster Johnson-Lindenstrauss projections.




Typically using sparse and structured matrices.

Next class: We will describe a construction where ΠA can be computed in $O(nd \log n)$ time.

RETURN TO SINGLE VECTOR PROBLEM

Goal: Develop methods that reduce a vector $\mathbf{x} \in \mathbb{R}^n$ down to $m \approx \frac{\log(1/\delta)}{\epsilon^2}$ dimensions in $o(mn)$ time and guarantee:

$$(1 - \epsilon)\|\mathbf{x}\|_2^2 \leq \|\mathbf{\Pi}\mathbf{x}\|_2^2 \leq (1 + \epsilon)\|\mathbf{x}\|_2^2$$



There is a truly brilliant method that runs in $O(n \log n)$ time.

Preview: Will involve Fast Fourier Transform in disguise.